

Prevalence of elongated styloid process on cone-beam computed tomographic images: a retrospective study

 Ezgi Katı

Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Hacettepe University, Ankara, Turkey

Cite this article: Katı E. Prevalence of elongated styloid process on cone-beam computed tomographic images: a retrospective study. *J Dent Sci Educ.* 2023;1(3):61-65.

Corresponding Author: Ezgi Katı, sahinkayaezgi@gmail.com

Received: 07/06/2023

Accepted: 26/06/2023

Published: 30/09/2023

ABSTRACT

Aims: The aim of the study was to evaluate the prevalence and the morphometric features of the styloid process (SP) via cone-beam computed tomography (CBCT).

Method: 400 SP of 200 individuals (92 males, 108 females, age range 12-81) were examined retrospectively in CBCT images in the axial, sagittal, and coronal planes. The measurements of SP and clinical and incidental findings associated with styloid ligament elongation (SPE) were detected. For descriptive statistics, the mean and standard deviation are given for numerical variables. The Mann-Whitney U test was used for measurements. The consistency of right and left SP lengths was analyzed for normality using the Shapiro-Wilk test ($p < 0.05$).

Results: Males (55.7%) had a substantially higher prevalence of ESP than females (33.1%) ($p = 0.032$). On the right side of the SP, the mean length was 26,76 mm (range: 8,7-63,3 mm), while on the left, it was 26,61 mm (range: 5,8-55,9 mm). 51 of them had bilateral lengths greater than 30 mm. The eight symptomatic patients' CBCT scans revealed SP sized anywhere from 36,6 mm to 69 mm on the right side. As incidental findings, soft tissue calcifications were detected in 5% of the 200 patients.

Conclusion: Men were found to have a higher prevalence of SPE than women, and a tiny percentage of cases were asymptomatic. There was no significant difference between right- and left-sided SP.

Key words: Calcification, cone-beam computed tomography, eagle syndrome

INTRODUCTION

The styloid process is an anatomical structure shaped like a cylindrical, thin, needle-like projection of varying lengths, averaging 2 to 3 cm.¹ It is classified as elongated if the measurement is more than 30 mm.² The styloid process (SP) projects from the inferior portion of the petrous temporal bone and offers attachment to the stylohyoid ligament and the muscles attached to the SP. Through these structures, SP facilitates the movement of the surrounding structures, such as the tongue, pharynx, larynx, hyoid bone, and mandible. Significant vessels and nerves surround the SP. The vena jugularis interna, arteria carotis interna, and some important cranial nerves lie medially to the SP. Arteria occipitalis and nervus hypoglossus run along its lateral side.³

The stylomastoid foramen, where the nervus facialis emerges from the skull, is directly posterior to the SP; however, it passes lateral to the process as it pierces through the parotid gland prior to splitting into its subsequent branches.^{4,5}

The structure shows variations in length, angulation, and other morphological features between individuals. The length of the styloid processes might also be different bilaterally within the same individual.⁶

Multiple theories have been laid out as the etiology responsible for the variance in ossification and elongation of SP. The first theory is the "theory of reactive hyperplasia". The second theory is the "theory of reactive metaplasia,". The third theory is the "theory of anatomic variance." An additional and fourth theory is that the elongated styloid process is due to retained embryologic tissue from Reichert's cartilage.⁷

Although these theories explain the ossification and elongations in different ways, a consensus has not been established.

The frequent presence of an elongated styloid process (SPE) is estimated at around 4% of the general population. Compared to men, women are more likely to have SPE.⁵

SPE may be a result of various clinical symptoms, such as pain in the neck and cervicofacial area. Although these physiological differences are often found incidentally, some patients might develop a constellation of symptoms known as Eagle syndrome. The symptomatology of Eagle syndrome occurs secondary to irritation and/or compression of surrounding structures from an abnormal SP.³

Despite the fact that the majority of these individuals have no symptoms, a small percentage of those with SPE show



symptoms and can present with one of two types of Eagle syndrome.^{7,8}

Eagle syndrome presents itself with various degrees of symptoms, such as a sharp pain in the neck or the ear that extends to the maxilla, face, and oral cavity. Moreover, it might also be correlated with vertigo, tinnitus, or a sensation of something abnormal in the pharynx. Symptoms of classic Eagle syndrome are usually unilateral but can rarely present bilaterally. Pressure on the arteria carotis interna might present with symptoms of internal carotid vascular insufficiency such as weakness, visual changes, or syncope exacerbated with head movement. The SPE might also pose the risk of carotid artery dissection, resulting in a stroke or transient ischemic attack.⁹

The diagnosis of Eagle syndrome depends on the patient's clinical presentation, radiological investigation, and lidocaine infiltration test.⁵

For radiological investigations, a lateral head and neck X-ray, a Towne radiograph, can identify the SPE in two-dimensional imaging. Computed tomography (CT) allows for the radiological examination and measurements of the SP three-dimensionally.^{10,11}

In the near past, three-dimensional cone-beam computerized tomography (CBCT) was a trustworthy imaging tool for anatomical structures of the maxillofacial region with high image quality, diagnostic accuracy, and sensitivity.^{12,13}

The purpose of the present study was to appraise the size of the styloid process, the prevalence of an elongated styloid ligament, its relation to sex, location, and correlation between the right and left sides using CBCT images.

METHODS

The radiographic images of 200 individuals who had been referred to the oral and maxillofacial radiology department for CBCT examination from September 2021 to August 2022 were evaluated retrospectively.

In this retrospective study, 200 CBCT images of patients who were randomly selected and aged from 12 to 81 years were analyzed. The study was carried out in the Department of Oral And Maxillofacial Radiology of Lokman Hekim University's Faculty of Dentistry. All the CBCT examinations were acquired with Newtom VGI-Evo (NewTom, Verona, Italy) and reconstructed with NNT® software (NewTom®, v. 5.6.0) with a 0.300 mm axial pitch and 0.300 mm axial thickness. The measurements were accomplished utilizing InVivo Dental Application Version 5.2.3 (Anatamage, Santa Clara, USA). Only high-quality images were included in the study. Any radiograph with questionable SP and images of low quality, such as scattering or insufficient accuracy of bony borders, was excluded from the study. The study protocol, numbered 2022/201, was approved by the Scientific Research Ethics Committee at Lokman Hekim University. The study's protocol was carried out according to the principles described in the Declaration of Helsinki, including all amendments and revisions. The assessment of images was fulfilled directly on the monitor screen (a 23-inch Acer 1920 x 1080 pixel HP Reconstruction monitor).

Radiological measurements of 400 SPs were performed on CBCT images of the 200 patients by an oral and maxillofacial radiologist. The length and type of SP were recorded separately for each side since there were some differences

between the right and left sides in the images. SP were assessed based on their average lengths. The length of the SP was recorded from the caudal margin on the tympanic plate to the tip of the process for each side. If the cranial part of the SP was not suitable for measurement, the length between the probable attachment point to the calvaria and the tip of the SP was recorded (14). The length of SP >30 mm was recorded as unilateral or bilateral elongation.^{2,5-21}

Statistical Analysis

The consistency of right and left styloid process lengths obtained from the subjects was analyzed for normality using the Shapiro-Wilk test. Neither of the measurements was found to be normally distributed among the study group. The presentation of descriptive studies used average ± standard deviation, and median presentation. The differences between the ages of the subjects according to gender were analyzed using the Student's t-test. A Mann-Whitney U test was used to analyze the difference between right and left styloid process length according to gender. A Kruskal-Wallis variance analysis was used to compare right and left styloid process length according to age groups. Spearman's rank correlation coefficient was calculated for the correlation between age and styloid process lengths. SPSS (version 21.00 for Windows. SPSS Inc., Chicago, IL, USA) was used for statistical analysis. (p <0.05)

RESULTS

A total of 200 individuals, 108 women and 92 men, between the ages of 12 and 81, with a mean age of 41,32±18,76 were selected for the research. (Table 1)

Table 1. Descriptive Analysis

	N	Minimum	Maximum	Mean	Std. Deviation
Age	200	12	81	41.32	18.769
Valid N (listwise)	200				

Examination of CBCT images demonstrated that the mean length of the styloid processes was 26,7 ± 8,3 mm (range: 8.7-63.3 mm) on the right side and 26,6 ± 7,7 mm (range: 5,8-55.9 mm) on the left side. The prevalence of ESP was significantly higher in males (55.7%) than females (33.1%). The consistency of right and left styloid process lengths obtained from the subjects was analyzed for normality using the Shapiro-Wilk test (p = 0.032). (Table 2).

Table 2. Test of Normality

Gender	Shapiro-Wilk		
	Statistic	df	Sig.
Female	.900	108	.32
Right ligament			
Male	.941	92	.169
Female	.842	108	.002
Left ligament			
Male	.931	92	.096

The results for the ligaments on each side were similar in both genders. (p=0.176 >0.05) (Table 3)

As incidental findings, 10 of 200 individuals had soft tissue calcifications. (Figure 1)

51 of the styloid processes were longer than 30 mm bilaterally. (Figure 2)



Table 3. Paired Samples Test for SP between right and left side			
	Mean	N	Std. Deviation
Right ligament	26.768	200	8.3125
Pair	26.613	200	7.7162
Left ligament			.176




Figure 1. CBCT Image of soft tissue calcifications. Tonsillolith, carotis artery calcifications, anthrolith.




Figure 2. Coronal, axial and sagittal views of CBCT image.



Only eight patients were symptomatic. The CBCT images of the symptomatic patients showed styloid processes measuring at least 36,6 mm to 69 mm on the right side. (Figure 3)




Figure 3. CBCT image of a SP measuring 69 mm.

DISCUSSION

There are several variations of features of the SP, such as the length, the degree of ossification, the distribution between genders, and the right and left sides.²²

The hyoid smaller horn, SP, and stylohyoid ligament are formed from Reichert's cartilage during the development of the head and neck region. In some cases, ossification of the stylohyoid ligament occurs and leads to the formation of SPE. The definition of Eagle's syndrome is symptomatic lengthening of the SP or calcification of the stylohyoid and stylomandibular ligaments.¹⁵

Physical examination and radiographic analysis are both used to make a diagnosis. There are several radiographic techniques used in dentistry to diagnose SPE, such as orthopantomographs and cone beam computed tomography.^{21,22,23-26}

In this study, CBCT images of 200 individuals were examined. Only a tiny part (between 4% and 10.3%) of those with SPEs, who represent roughly 4% of the population, are considered to show symptoms.²⁷

Öztunç et al.²³ compared elongated SP and without elongated SP on CBCT images and found the mean age of the patients with SPE to be 47.28 ± 12.4 years. In this study, the mean age of the patients with SPE was $41,32 \pm 18,76$ years (range: 12–81 years), and similar to this, all of the symptomatic patients were older than .⁴⁰

Similar to this study, other studies have found that the mean SP length was significantly different between males and females. The prevalence of SPE was higher in males than females.^{21,25,28}

Camarda et al.²⁹ measured the mean length of the SP in their study, and it was similar to this study. And also, the values of both sides were found to be close to each other, similar to the present study. There was no difference between both sides in both genders in the present study.

Jung et al.³⁰ examined panoramic images of 1000 patients and reported that both sides' distributions are similar and do not significantly differ (Kolmogorov–Smirnov test, $P= 0.380$).

Although the examined image technique differs between 2 and 3 dimensions, both studies showed similar results.

Clinical complaints of SPE are related to the size of the stylohyoid complex.¹⁹ In this study, the CBCT images of the eight symptomatic cases showed SP measuring at least 36,6 mm to 69 mm on the right side. Camarda et al.¹⁹ reported that patients experiencing symptoms tended to be older than 40 and had no recent history of cervicopharyngeal trauma, tonsillectomy, or other surgery. Nobody among our symptomatic individuals had ever undergone surgery or experienced trauma.

Soft tissue calcifications such as tonsilloliths, antroliths, and carotid artery calcifications were detected as incidental findings.

Missias et al.³¹ detected soft tissue calcifications in CBCT images with different fields of view. SPE and tonsillolith were the most frequent calcifications in their study. Pette et al.³² evaluated the prevalence of soft tissue calcifications in 318 samples and reported that tonsilloliths and antroliths were common with a range of 32% and 3%, respectively. Price et al.³³ evaluated incidental findings in the maxillofacial region by CBCT and reported tonsilloliths, antroliths, and carotid artery calcifications at 43%, 5%, and 13% in CBCT scans of 300 individuals, respectively.

In the present study, 10 of 200 individuals had soft tissue calcifications. Tonsilloliths (4 of 10 individuals) located on the right side of the lateral wall of the pharynx were sized at 0.8mm, 1,8 mm, 1,4 mm, 1,3 mm, and 0,7 mm in diameter. Soft tissue calcifications (3 of 10 individuals) located in the laryngopharynx sized 1,7 mm, 0,8 mm, and 0,5 mm in diameter. Antrolith (1 of 10) sized 2 mm in diameter and mucosal thickening consistent with maxillary sinusitis. The elongated styloid process might also compress the carotid artery. In this study, a CBCT image showed carotid artery calcifications (2 of 10 individuals) sized 2,0 and 2,2 mm in diameter.

CONCLUSION

In this study, morphometric examination of the styloid process, clinical complaints, and potential incidental findings were all assessed using CBCT images. While the prevalence of SPE was found to be higher in men compared to the female population, no significant difference was found for right- and left-sided SP.

ETHICAL DECLARATIONS

Ethics Committee Approval: The study protocol, numbered 2022/201, was approved by the Scientific Research Ethics Committee at Lokman Hekim University.

Informed Consent: Because the study was designed retrospectively, no written informed consent form was obtained from patients.

Referee Evaluation Process: Externally peer-reviewed.

Conflict of Interest Statement: The authors have no conflicts of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Author Contributions: All of the authors declare that they have all participated in the design, execution, and analysis of the paper, and that they have approved the final version.



REFERENCES

- Glogoff MR, Baum SM, Cheifetz I. Diagnosis and treatment of Eagle's Syndrome. *J Oral Surg*. 1981;39(12):941-944.
- Murtagh RD, Caracciolo JT, Fernandez G. CT findings associated with Eagle syndrome. *Am J Neuroradiol*. 2001;22(7):1401-1412.
- Abuhaimeed AK, Alvarez R, Menezes RG. Anatomy, Head and Neck, Styloid Process. 2022 Jan 14. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. PMID: 31082019.
- Vadgaonkar R, Murlimanju BV, Prabhu LV, et al. Morphological study of styloid process of the temporal bone and its clinical implications. *Anat Cell Biol*. 2015;48(3):195-200.
- Piagkou M, Anagnostopoulou S, Kouladouros K, Piagkos G. Eagle's syndrome: a review of the literature. *Clin Anat*. 2009;22(5):545-558.
- Custodio AL, Silva MR, Abreu MH, Araújo LR, de Oliveira LJ. Styloid process of the temporal bone: morphometric analysis and clinical implications. *Biomed Res Int*. 2016;2016:8792725.
- Badhey A, Jategaonkar A, Anglin Kovacs AJ, Kadakia S, De Deyn PP, Ducic Y, et al. Eagle syndrome: a comprehensive review. *Clin Neurol Neurosurg*. 2017;159:34-38.
- Moon CS, Lee BS, Kwon YD, et al. Eagle's syndrome: a case report. *J Korean Assoc Oral Maxillofac Surg*. 2014;40(1):43-47.
- Kamal A, Nazir R, Usman M, Salam BU, Sana F. Eagle syndrome; radiological evaluation and management. *J Pak Med Assoc*. 2014;64(11):1315-1317.
- Nayak DR, Pujary K, Aggarwal M, Punnoose SE, Chaly VA. Role of three-dimensional computed tomography reconstruction in the management of elongated styloid process: a preliminary study. *J Laryngol Otol*. 2007;121(4):349-353.
- Petrović B, Radak D, Kostić V, Covicković-Sternić N. Styloid syndrome: a review of literature. *Srp Arh Celok Lek*. 2008;136(11-12):667-674.
- Scarfe WC, Farman AG, Sukovic P. Clinical applications of cone-beam computed tomography in dental practice. *J Canadian Dental Assoc*. 2006;72(1):75-80.
- White SC, Pharoah MJ. *White and Pharoah's Oral Radiology: Principles and Interpretation*. 8th ed. St. Louis, Missouri: Elsevier Mosby; 2018. 185-199.
- Paparella MM, Shumrick DA. *Otolaryngology Philadelphia*. 2nd ed. Saunders: Elsevier; 1980.
- Correll RW, Jensen JL, Taylor JB, Rhyne RR. Mineralization of the stylohyoid-stylomandibular ligament complex. A radiographic incidence study. *Oral Surg Oral Med Oral Pathol*. 1979;48(4):286-291.
- Keur JJ, Campbell JP, McCarthy JF, Ralph WJ. The clinical significance of the elongated styloid process. *Oral Surg Oral Med Oral Pathol*. 1986;61(4):399-404.
- Langlais RP, Miles DA, Van Dis ML. Elongated and mineralized stylohyoid ligament complex: a proposed classification and report of a case of Eagle's syndrome. *Oral Surg Oral Med Oral Pathol*. 1986;61(5):527-532.
- Sekerci AE, Soylu E, Arikan MP, Aglarci OS. Is there a relationship between the presence of ponticulus posticus and elongated styloid process? *Clin Imaging*. 2015;39(2):220-224.
- Anbiaee N, Javadzadeh A. Elongated styloid process: Is it a pathologic condition? *Indian J Dent Res*. 2011;22(5):673-677.
- Prasad KC, Kamath MP, Reddy KJ, Raju K, Agarwal S. Elongated styloid process (Eagle's syndrome): A clinical study. *J Oral Maxillofac Surg*. 2002;60(2):171-175.
- Donmez M, Okumus O, Pekiner FN. Cone beam computed tomographic evaluation of styloid process: A retrospective study of 1000 patients. *European J Dentistry*. 2017;11(02):210-215.
- İlgü M, İlgü D, Güler N, Bayırlı G. Incidence of the type and calcification patterns in patients with elongated styloid process. *J Int Med Res*. 2005;33(1):96-102.
- Öztunç H, Evlice B, Tatlı U, Evlice A. Cone-beam computed tomographic evaluation of styloid process: a retrospective study of 208 patients with orofacial pain. *Head & Face Med*. 2014;10(1):1-7.
- Tijanić M, Burić N, Burić K. The use of Cone Beam CT (CBCT) in differentiation of true from mimicking Eagle's syndrome. *Int J Environ Res Public Health*. 2020;17(16):5654.
- Buyuk C, Gunduz K, Avsever H. Morphological assessment of the stylohyoid complex variations with cone beam computed tomography in a Turkish population. *Folia morphologica* 2018;77(1):79-89.
- Javadian Langaroodi A, Hoseini Zarch SH, Rahpeyma A, Ebrahimnejad H, Arezoobakhsh A, Sanaei A. Assessment of stylohyoid ligament in patients with Eagle's syndrome and patients with asymptomatic elongated styloid process: A cone-beam computed tomography study. *Journal of Oral Health and Oral Epidemiology*. 2016;5(4):215-220.
- Rechtweg JS, Wax MK. Eagle's syndrome: a review. *Am J Otolaryngol*. 1998;19(5):316-321.
- Nalçacı R, Mısırlıoğlu M. Yaşlı bireylerde stiloid sürecin radyolojik olarak değerlendirilmesi. *AÜ Dış Hekimliği Fakültesi Derg*. 2006;3(1):1-6.
- Camarda AJ, Deschamps C, Forest DI. Stylohyoid chain ossification: a discussion of etiology. *Oral Surg Oral Med Oral Pathol*. 1989;67(5):508-514.
- Jung T, Tschernitschek H, Hippen H, Schneider B, Borchers L. Elongated styloid process: when is it really elongated?. *Dentomaxillofacial Radiology*. 2004;33(2):119-124.
- Missias EM, Nascimento EHL, Pontual MLA, Pontual AA, Freitas DQ, Perez DEC, Ramos-Perez FMM. Prevalence of soft tissue calcifications in the maxillofacial region detected by cone beam CT. *Oral Diseases*. 2018;24(4):628-637.
- Pette GA, Norkin FJ, Ganeles J, et al. Incidental findings from a retrospective study of 318 cone beam computed tomography consultation reports. *Int J Oral Maxillofac Implants*. 2012;27(3):595-603.
- Price JB, Thaw KL, Tyndall DA, Ludlow JB, Padilla RJ. Incidental findings from cone beam computed tomography of the maxillofacial region: a descriptive retrospective study. *Clin Oral Implants Res*. 2012;23(11):1261-1268.